Week 2
Term 1
2022

HAWKER COLLEGE
Engage | Inspire | Achieve

MM3
Further differentiation and applications

Goals

By the end of this week you will:

- understand the notion of composition of functions and use the chain rule for determining the derivatives of composite functions
- estimate the limit of $\frac{a^h-1}{h}$ as $h \to 0$ using technology, for various values of $a > 0$
- recognise that e is the unique number a for which the above limit is 1
- establish and use the formula $\frac{d}{dx}(e^x) = e^x$
- use exponential functions and their derivatives to solve practical problems.

Theoretical Components

Resources:

- *Year 12 Maths Quest Methods* Chapter 7

Chain Rule

$$\frac{dy}{dx} = \frac{dy}{du} \times \frac{du}{dx}$$

https://www.khanacademy.org/math/ap-calculus-ab/product-quotient-chain-rules-ab/chain-rule-proof-ab/v/chain-rule-proof

Read and make notes on examples 13, 14 and 15 from Chapter 7

What is base e?
Read through to get an insight on the number 'e':
http://bit.ly/w8OiD

http://www.mathopolis.com/questions/q.php?id=2011&site=1&ref=/numbers/e-eulers-number.html&qs=2011_2012_2013

Derivative of $y = e^x$, and $y = e^{f(x)}$ from first principles
See the following page

Read and make notes on examples 16 - 19 from Chapter 7

Practical Components

Complete the following questions. Organise your solutions neatly in your exercise book.

You will require Chapter 7 of Maths Quest Methods (pdf – Google Classroom).

Ex 7D The Chain Rule
Q's 1, 2, 4, 5, 6 , 7 (b,d,f,h), 8, 10 (c,d), 13

Ex 7E The Derivative of e^x and $e^{f(x)}$
Q's 1 (a,d,g), 2 (b,e,h,k), 3, 4 (a,d,g), 5 (c,f,i,l), 6, 7, 8, 9

Investigation

See separate page

QFO

Quiz/Forum/Other

Remember to check hawkermaths.com for each week's learning brief, and check in with your teacher each lesson.
No Mathspace this week ☺

Chain rule

1. A composite function is a function composed of two (or more) functions.
2. Composite functions can be differentiated using the chain rule, $\frac{dy}{dx} = \frac{dy}{du} \times \frac{du}{dx}$.
3. A short way of applying the chain rule is:
   If $f(x) = [g(x)]^n$ then $f'(x) = n[g(x)]^{n-1} \times g'(x)$.

The derivative of e^x

1. If $f(x) = e^x, f'(x) = e^x$.
2. If $f(x) = e^{kx}, f'(x) = ke^{kx}$.
3. If $f(x) = ae^{kx+c}, f'(x) = ake^{kx+c}$.
4. If $f(x) = ae^{g(x)}, f'(x) = g'(x) \times ae^{g(x)}$.

Derivative of $f(x) = e^x$ from first principles:

$$f'(x) = \lim_{h \to 0} \frac{f(x+h) - f(x)}{h}, h \neq 0$$

$$= \lim_{h \to 0} \frac{e^{x+h} - e^x}{h}$$

$$= \lim_{h \to 0} \frac{e^x e^h - e^x}{h}$$

$$= \lim_{h \to 0} \frac{e^x(e^h - 1)}{h}$$

$$= e^x \lim_{h \to 0} \frac{e^h - 1}{h}$$

Note that $\lim_{h \to 0} \frac{e^h - 1}{h}$ can be deduced by using a calculator and substituting values of h close to zero.

h	$\frac{e^h - 1}{h}$
0.01	1.0050
0.0001	1.000 05
0.000 001	1.000 000

That is, $\lim_{h \to 0} \frac{e^h - 1}{h} = 1$.

Therefore, $f'(x) = e^x \times 1$
$= e^x$

If $f(x) = e^x$ then $f'(x) = e^x$.

Note: e^x is the only function which has itself as a derivative.

MM3 Week 2 Investigation

The graph of $y = Ne^{kx}$

The diagram at right shows the graph of $y = e^x$ and uses the letters A, B and C to indicate key parts of the graph.



In this investigation you will use your graphics calculator to observe and report on the effect of changing N and k in the equation $y = Ne^{kx}$.

1 $N = 2$. On the same axes, graph the equations $y = 2e^x$ and $y = e^x$.
In your book, sketch the view window.
Write a sentence summarising the effect of changing N from 1 to 2.

2 $N = -1$. On the same axes, graph the equations $y = -1 \times e^x$ and $y = e^x$.
In your book, sketch the view window.
Write a sentence summarising the effect of changing N from 1 to -1.

3 $k < 0$. On the same axes, graph the equations $y = e^x$ and $y = e^{-x}$.
In your book, sketch the view window.
Write a sentence summarising the effect of changing k from 1 to -1.

4 $0 < k < 1$. On the same axes, graph the equations $y = e^x$ and $y = e^{0.5x}$.
In your book, sketch the view window.
Write a sentence summarising the effect of changing k from 1 to 0.5.

5 $k > 1$. On the same axes, graph the equations $y = e^x$ and $y = e^{2x}$.
In your book, sketch the view window.
Write a sentence summarising the effect of changing k from 1 to 2.

Challenge

Use your calculator to obtain a guess-and-check solution to the following problem. Find the values of k and N such that the graph of $y = Ne^{kx}$ passes through $(-2, 4)$ and $(0, 2)$, the points shown.



Write down your guesses and the adjustments you made to find the equation of the challenge question.

Complete the challenge algebraically as well.

You can use https://www.desmos.com/calculator or CAS calculator for your graphs.